

DATA SHEET



Klentaq DNA Polymerase

Thermostable DNA polymerase
Thermus aquaticus, recombinant, *E. coli*

Cat. Nº.	Amount
<input type="checkbox"/> POL-127XS	250 units
<input type="checkbox"/> POL-127S	500 units
<input type="checkbox"/> POL-127M	1.000 units
<input type="checkbox"/> POL-127L	2 x 1.000 units (M)
<input type="checkbox"/> POL-127XL	4 x 1.000 units (M)

Unit Definition:

One unit is defined as the amount of the enzyme required to catalyze the incorporation of 10 nmoles of dNTPs into an acid-insoluble form in 30 minutes at 74 °C.

Concentration:

5 units/μL

Shipping:

Shipped on blue ice

Storage Conditions:

Store at -20 °C

For *in vitro* use only!

Additional Storage Conditions:

Avoid freeze/thaw cycles

Shelf Life:

12 months

Kit contents:

Klentaq (blue cap)

5 units/μL Klentaq DNA Polymerase in Tris-HCl pH 8.0 (25 °C), KCl, EDTA, DTT, 50% (v/v) Glycerol and stabilizers.

Klentaq Reaction Buffer complete (red cap) - 10x conc.

Tris-HCl pH 8.5 (25°C), KCl and 25 mM MgCl₂.

Description:

Klentaq is a 5'-exonuclease deficient Taq polymerase (an N-terminal deletion of Taq) with improved fidelity and thermostability.

PCR Reaction Setup

The PCR procedure below shows appropriate volumes for a single 50 μL reaction. For multiple reactions, prepare a master mix of components common to all and then dispense appropriate volumes into each PCR reaction tube prior to adding template DNA and primers.

Thaw, mix, and briefly centrifuge each component before use. Add the following components to a microcentrifuge tube:

1. Prepare PCR master mix

Note: Consider the volumes for all components listed next steps to determine the correct amount of water required to reach your final reaction volume.

Components	50 μL rxn	[final]
Water, grade PCR	To 50 μl	
10x Reaction Buffer	5 μl	1X
dNTP (Mix 10 mM)	1 μl	200 μM
Klentaq (5 U/ μl)	0,5 μl	2,5 U/reaction

Mix and briefly centrifuge the components.

2. Add template DNA and primers

Components	50 μL rxn	[final]
Foward primer (10 μM)	0,5 - 2,5 μl	0,1 - 0,5 μM
Reverse primer (10 μM)	0,5 - 2,5 μl	0,1 - 0,5 μM
DNA template		10 pg - 1 μg**

**genomic DNA: 1 ng-1μg; plasmidial or viral DNA: 1 pg-1 ng

Cap each tube, mix, and briefly centrifuge the content.

3. Incubate reactions in a thermal cycler.

Recommended cycling conditions:

Step	Temp.	Time
Initial denaturation	95 °C	2 min
25 - 40 cycles	Denaturation	95 °C
	Annealing ¹	45-68 °C
	Elongation ²	68 °C
Final extension (optional)	68 °C	1 - 2 min
Hold	4 - 8 °C	

1)The annealing temperature depends on the melting temperature of the primers used.

2)The elongation time depends on the length of the fragments to be amplified. A time of 2 min/kb is recommended.

For optimal specificity and amplification an individual optimization of the recommended parameters may be necessary for each new template DNA and/or primer pair.