




5-Methyl-dCDP

5-Methyl-2'-deoxycytidine-5'-diphosphate, Sodium salt

Cat. No.	Amount
NU-1163S	50 µl (10 mM)
NU-1163L	5 x 50 µl (10 mM)



Structural formula of 5-Methyl-dCDP

For general laboratory use.

Shipping: shipped on gel packs

Storage Conditions: store at -20 °C

Short term exposure (up to 1 week cumulative) to ambient temperature possible.

Shelf Life: 12 months after date of delivery

Molecular Formula: C₁₀H₁₇N₃O₁₀P₂ (free acid)

Molecular Weight: 401.20 g/mol (free acid)

Exact Mass: 401.04 g/mol (free acid)

CAS#: 22003-12-9

Purity: ≥ 95 % (HPLC)

Form: solution in water

Color: colorless to slightly yellow

Concentration: 10 mM - 11 mM

pH: 7.5 ±0.5

Spectroscopic Properties: λ_{max} 277 nm, ε 9.0 L mmol⁻¹ cm⁻¹ (Tris-HCl pH 7.5)

Selected References:

Kamiya *et al.* (2004) Important amino acids in the phosphohydrolase module of *Escherichia coli* Orf135. *Biochem Biophys Res Commun.* **323** (3):1063.

Holliday *et al.* (2002) DNA methylation and epigenetic inheritance. *Methods.* **27** (2):179.

Kaito *et al.* (2001) Activation of the maternally preset program of apoptosis by microinjection of 5-aza-2'-deoxycytidine and 5-methyl-2'-deoxycytidine-5'-triphosphate in *Xenopus laevis* embryos. *Dev Growth Differ.* **43** (4):383.

Wong *et al.* (1997) A novel method for producing partial restriction digestion of DNA fragments by PCR with 5-methyl-CTP. *Nucleic Acids Res.* **25** (20):4169.

Chen *et al.* (1993) Direct induction of DNA hypermethylation in sea urchin embryos by microinjection of 5-methyl dCTP stimulates early histone gene expression and leads to developmental arrest. *Dev Biol.* **155** (1):75.